

Références

Polyarthrite rhumatoïde, auto-antigène(s) et auto-anticorps : état des lieux et actualités récentes

Eléonore Bettacchioli, Laboratoire d'immunologie du CHU de Brest.

- Smolen JS, Aletaha D, McInnes IB. Rheumatoid arthritis. *The Lancet*. 22 oct 2016;388(10055):2023-38.
- Rheumatoid arthritis. *Nat Rev Dis Primer*. 8 févr 2018;4:18002.
- McInnes IB, Schett G. The Pathogenesis of Rheumatoid Arthritis. *N Engl J Med*. 8 déc 2011;365(23):2205-19.
- Aletaha D, Neogi T, Silman AJ, Funovits J, Felson DT, Bingham CO, et al. 2010 Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis Rheum*. sept 2010;62(9):2569-81.
- Haro I, Sanmartí R, Gomara MJ. Implications of Post-Translational Modifications in Autoimmunity with Emphasis on Citrullination, Homocitrullination and Acetylation for the Pathogenesis, Diagnosis and Prognosis of Rheumatoid Arthritis. *Int J Mol Sci*. 2022 Dec 13;23(24):15803.
- De Stefano L, D'Onofrio B, Manzo A, Montecucco C, Bugatti S. The Genetic, Environmental, and Immunopathological Complexity of Autoantibody-Negative Rheumatoid Arthritis. *Int J Mol Sci*. 2021 Nov 17;22(22):12386.
- Girbal-Neuhausser E, Durieux JJ, Arnaud M, Dalbon P, Sebbag M, Vincent C, et al. The epitopes targeted by the rheumatoid arthritis-associated antifilaggrin autoantibodies are posttranslationally generated on various sites of (pro)filaggrin by deimination of arginine residues. *J Immunol Baltim Md 1950*. 1 janv 1999;162(1):585-94.
- Trier NH, Houen G. Epitope Specificity of Anti-Citrullinated Protein Antibodies. *Antibodies*. mars 2017;6(1):5.
- Brentville VA, Vankemmelbeke M, Metheringham RL, Durrant LG. Post-translational modifications such as citrullination are excellent targets for cancer therapy. *Semin Immunol*. 1 févr 2020;47:101393.
- Hill JA, Southwood S, Sette A, Jevnikar AM, Bell DA, Cairns E. Cutting Edge: The Conversion of Arginine to Citrulline Allows for a High-Affinity Peptide Interaction with the Rheumatoid Arthritis-Associated HLA-DRB1*0401 MHC Class II Molecule. *J Immunol*. 15 juill 2003;171(2):538-41.
- Desclos-Theveniau M, Bonnaure-Mallet M, Meuric V. Peptidylarginine désiminasés du microbiote buccal et polyarthrite rhumatoïde [Protein arginine deiminase of oral microbiome plays a causal role in the polyarthritis rheumatoid initiating]. *Med Sci (Paris)*. 2020 May;36(5):465-471.
- Wegner N, Wait R, Sroka A, Eick S, Nguyen K-A, Lundberg K, et al. Peptidylarginine deiminase from *Porphyromonas gingivalis* citrullinates human fibrinogen and alpha-enolase: implications for autoimmunity in rheumatoid arthritis. *Arthritis Rheum*. sept 2010;62(9):2662-72.
- Martinez-Prat L, Nissen MJ, Lamacchia C, Bentow C, Cesana L, Roux-Lombard P, et al. Comparison of Serological Biomarkers in Rheumatoid Arthritis and Their Combination to Improve Diagnostic Performance. *Front Immunol*. 2018;9:1113.
- Klareskog L, Stolt P, Lundberg K, Källberg H, Bengtsson C, Grunewald J, et al. A new model for an etiology of rheumatoid arthritis: smoking may trigger HLA-DR (shared epitope)-restricted immune reactions to autoantigens modified by citrullination. *Arthritis Rheum*. janv 2006;54(1):38-46.
- Humphreys JH, van Nies JA, Chipping J, Marshall T, van der Helm-van Mil AH, Symmons DP, Verstappen SM. Rheumatoid factor and anti-citrullinated protein antibody positivity, but not level, are associated with increased mortality in patients with rheumatoid arthritis: results from two large independent cohorts. *Arthritis Res Ther*. 2014 Dec 4;16(6):483.

- Bugatti S, Bogliolo L, Vitolo B, Manzo A, Montecucco C, Caporali R. Anti-citrullinated protein antibodies and high levels of rheumatoid factor are associated with systemic bone loss in patients with early untreated rheumatoid arthritis. *Arthritis Res Ther*. 2016 Oct 6;18(1):226.
- Reynisdottir G, Karimi R, Joshua V, Olsen H, Hensvold AH, Harju A, Engström M, Grunewald J, Nyren S, Eklund A, Klareskog L, Sköld CM, Catrina AI. Structural changes and antibody enrichment in the lungs are early features of anti-citrullinated protein antibody-positive rheumatoid arthritis. *Arthritis Rheumatol*. 2014 Jan;66(1):31-9.
- Alemao E, Guo Z, Frits ML, Iannaccone CK, Shadick NA, Weinblatt ME. Association of anti-cyclic citrullinated protein antibodies, erosions, and rheumatoid factor with disease activity and work productivity: A patient registry study. *Semin Arthritis Rheum*. 2018 Apr;47(5):630-638.
- Cantagrel A, Degboé Y. Les nouveaux auto-anticorps de la polyarthrite rhumatoïde : des anticorps qui reconnaissent des protéines du soi ayant subi des modifications post-translationnelles. *Rev Rhum*. 1 janv 2016;83(1):12-8.
- Mydel P, Wang Z, Brisslert M, Hellvard A, Dahlberg LE, Hazen SL, et al. Carbamylation-dependent activation of T cells: a novel mechanism in the pathogenesis of autoimmune arthritis. *J Immunol Baltim Md 1950*. 15 juin 2010;184(12):6882-90.
- Jiang X, Trouw LA, van Wesemael TJ, Shi J, Bengtsson C, Källberg H, et al. Anti-CarP antibodies in two large cohorts of patients with rheumatoid arthritis and their relationship to genetic risk factors, cigarette smoking and other autoantibodies. *Ann Rheum Dis*. oct 2014;73(10):1761-8.
- Shi J, van de Stadt LA, Levarht EWN, Huizinga TWJ, Hamann D, van Schaardenburg D, et al. Anti-carbamylated protein (anti-CarP) antibodies precede the onset of rheumatoid arthritis. *Ann Rheum Dis*. avr 2014;73(4):780-3.
- Griffiths HR. Is the generation of neo-antigenic determinants by free radicals central to the development of autoimmune rheumatoid disease? *Autoimmun Rev*. juill 2008;7(7):544-9.
- Nissim A, Winyard PG, Corrigan V, Fatah R, Perrett D, Panayi G, et al. Generation of neoantigenic epitopes after posttranslational modification of type II collagen by factors present within the inflamed joint. *Arthritis Rheum*. déc 2005;52(12):3829-38.
- Strollo R, Ponchel F, Malmström V, Rizzo P, Bombardieri M, Wenham CY, et al. Autoantibodies to posttranslationally modified type II collagen as potential biomarkers for rheumatoid arthritis. *Arthritis Rheum*. juill 2013;65(7):1702-12.
- He Y, Ge C, Moreno-Giró À, Xu B, Beusch CM, Sandor K, et al. A subset of antibodies targeting citrullinated proteins confers protection from rheumatoid arthritis. *Nat Commun*. 2023 Feb 8;14(1):691.
- Raposo B, Afonso M, Israelsson L, Wähämaa H, Stålesen R, Wermeling F, et al. Divergent and dominant anti-inflammatory effects of patient-derived anticitrullinated protein antibodies (ACPA) in arthritis development. *Ann Rheum Dis*. 2023 May;82(5):724-726.